

Literatur Themenheft Demenz

„Ärzteblatt Sachsen“, Heft 8/2011

Autorin: V.A. Holthoff

Soziotherapeutische Behandlungsmöglichkeiten bei Demenzen

- AAGP, A. (2003) Consensus statement on improving the quality of mental health care in U.S. nursing homes: management of depression and behavioral symptoms associated with dementia. In J Am Geriatr Soc, pp. 1287-1298.
- Baillon, S., Van Diepen, E., Prettyman, R., et al (2004) A comparison of the effects of Snoezelen and reminiscence therapy on the agitated behaviour of patients with dementia. Int J Geriatr Psychiatry, 19, 1047-1052.
- Ballard, C., Brown, R., Fossey, J., et al (2009) Brief psychosocial therapy for the treatment of agitation in Alzheimer disease (the CALM-AD trial). Am J Geriatr Psychiatry, 17, 726-733.
- Bienstein, C. F., A. (2003) Basale Stimulation in der Pflege. Seelzer/Velber: Kallmeyer Verlag.
- BMG (2006) Bundesministerium für Gesundheit Rahmenempfehlungen zum Umgang mit herausforderndem Verhalten bei Menschen mit Demenz in der stationären Altenhilfe pp.
www.bmg.bund.de/.../Forschungsberichte/f007,templateId=raw,property=publicationFile.pdf/f007.pdf
- Brodaty, H., Green, A. & Koschera, A. (2003) Meta-analysis of psychosocial interventions for caregivers of people with dementia. J Am Geriatr Soc, 51, 657-664.
- Chung, J. C., Lai, C. K., Chung, P. M., et al (2002) Snoezelen for dementia. Cochrane Database Syst Rev, CD003152.
- Clare, L., Woods, R. T., Moniz Cook, E. D., et al (2003) Cognitive rehabilitation and cognitive training for early-stage Alzheimer's disease and vascular dementia. Cochrane Database Syst Rev, CD003260.
- Cohen-Mansfield, J. (2001) Nonpharmacologic interventions for inappropriate behaviors in dementia: a review, summary, and critique. Am J Geriatr Psychiatry, 9, 361-381.
- Eggermont, L. H., Swaab, D. F., Hol, E. M., et al (2009) Walking the line: a randomised trial on the effects of a short term walking programme on cognition in dementia. J Neurol Neurosurg Psychiatry, 80, 802-804.
- Feil, N. (1993) Validation. Ein neuer Weg zum Verständnis alter Menschen. ISBN 3-90-129102-4
- Feil, N., Sutton, E., Johnson, F. (2001): Trainingsprogramm Validation. ISBN 3-49-701558-X
- Filan, S. L. & Llewellyn-Jones, R. H. (2006) Animal-assisted therapy for dementia: a review of the literature. Int Psychogeriatr, 18, 597-611.
- Forbes, D., Forbes, S., Morgan, D. G., et al (2008) Physical activity programs for persons with dementia. Cochrane Database Syst Rev, CD006489.
- Forbes, D., Morgan, D. G., Bangma, J., et al (2004) Light therapy for managing sleep, behaviour, and mood disturbances in dementia. Cochrane Database Syst Rev, CD003946.
- Gitlin, L. N., Winter, L., Burke, J., et al (2008) Tailored activities to manage neuropsychiatric behaviors in persons with dementia and reduce caregiver burden: a randomized pilot study. Am J Geriatr Psychiatry, 16, 229-239.
- Gitlin, L. N., Winter, L., Dennis, M. P., et al (2010) Targeting and managing behavioral symptoms in individuals with dementia: a randomized trial of a nonpharmacological intervention. J Am Geriatr Soc, 58, 1465-1474.

- Gitlin, L. N., Winter, L., Vause Earland, T., et al (2009) The Tailored Activity Program to reduce behavioral symptoms in individuals with dementia: feasibility, acceptability, and replication potential. *Gerontologist*, 49, 428-439.
- Graff, M. J., Vernooij-Dassen, M. J., Thijssen, M., et al (2007) Effects of community occupational therapy on quality of life, mood, and health status in dementia patients and their caregivers: a randomized controlled trial. *J Gerontol A Biol Sci Med Sci*, 62, 1002-1009.
- Graff, M. J., Vernooij-Dassen, M. J., Thijssen, M., et al (2006) Community based occupational therapy for patients with dementia and their care givers: randomised controlled trial. *BMJ*, 333, 1196.
- Guetin, S., Portet, F., Picot, M. C., et al (2009) Effect of music therapy on anxiety and depression in patients with Alzheimer's type dementia: randomised, controlled study. *Dement Geriatr Cogn Disord*, 28, 36-46.
- Hermans, D. G., Htay, U. H. & McShane, R. (2007) Non-pharmacological interventions for wandering of people with dementia in the domestic setting. *Cochrane Database Syst Rev*, CD005994.
- Heyn, P., Abreu, B. C. & Ottenbacher, K. J. (2004) The effects of exercise training on elderly persons with cognitive impairment and dementia: a meta-analysis. *Arch Phys Med Rehabil*, 85, 1694-1704.
- Holthoff, V., Marschner, K., Gerner, A., Juranz, L., Schützwohl, M., Kallert, L., Becker, T. Reuster, T. (2011) ERGODEM-Individualisierte Ergotherapie in der Demenzbehandlung ist effektiv 10. Kongress der Deutschen Gesellschaft für Gerontopsychiatrie und -psychotherapie e.V. 11.-13. Mai 2011 in Berlin
- IQWiG (2009) Nichtmedikamentöse Behandlung der Alzheimer Demenz.1. Version des Abschlussberichtes im Auftrag des Gemeinsamen Bundesausschusses.
- Kemoun, G., Thibaud, M., Roumagne, N., et al (2010) Effects of a physical training programme on cognitive function and walking efficiency in elderly persons with dementia. *Dement Geriatr Cogn Disord*, 29, 109-114.
- Law M, Polatajko H, Carswell A, McColl MA, Pollock N, Baptiste S (1999). Das Kanadische Modell der Occupational Performanz und das Canadian Occupational Performanz Measure. In: Jerosch-Herold C, Marotzki U, Back BM, Weber P (eds.), *Ergotherapie - Reflexion und Analyse. Konzeptionelle Modelle für die ergotherapeutische Praxis*. 1. Aufl. Berlin: Springer, 156-173
- Livingston, G., Johnston, K., Katona, C., et al (2005) Systematic review of psychological approaches to the management of neuropsychiatric symptoms of dementia. *Am J Psychiatry*, 162, 1996-2021.
- Marquardt, G. (2007) Kriterienkatalog Demenzfreundliche Architektur, <http://nbn-resolving.de/urn:nbn:de:swb:14-1185359165306-51589>
- Neal, M. & Briggs, M. (2003) Validation therapy for dementia. *Cochrane Database Syst Rev*, CD001394.
- Oswald, W. D., Hagen, B. & Rupprecht, R. (2001) [Non-medicamentous therapy and prevention of Alzheimer disease]. *Z Gerontol Geriatr*, 34, 116-121.
- Perkins, J., Bartlett, H., Travers, C., et al (2008) Dog-assisted therapy for older people with dementia: a review. *Australas J Ageing*, 27, 177-182.
- Raglio, A., Oasi, O., Gianotti, M., et al (2010) Effects of music therapy on psychological symptoms and heart rate variability in patients with dementia. A pilot study. *Curr Aging Sci*, 3, 242-246.
- Reuster, T. J., L. Schützwohl, M. Holthoff, V. (2008) Effektivität einer optimierten Ergotherapie bei Demenz im häuslichen Setting (ERGODEM). *Zeitschrift für Gerontopsychologie & -psychiatrie*, 21, 185-189.
- Robinson, L., Hutchings, D., Corner, L., et al (2006) A systematic literature review of the effectiveness of non-pharmacological interventions to prevent wandering in dementia and evaluation of the ethical implications and acceptability of their use. *Health Technol Assess*, 10, iii, ix-108.
- Rolland, Y., Pillard, F., Klapouszczak, A., et al (2007) Exercise program for nursing home residents with Alzheimer's disease: a 1-year randomized, controlled trial. *J Am Geriatr Soc*, 55, 158-165.

- Romero, B. & Wenz, M. (2002) [Concept and effectiveness of a treatment program for patients with dementia and their relatives. Results from the Bad Aibling Alzheimer Disease Therapy Center]. *Z Gerontol Geriatr*, 35, 118-128.
- Schwenk, M., Zieschang, T., Oster, P., et al (2010) Dual-task performances can be improved in patients with dementia: a randomized controlled trial. *Neurology*, 74, 1961-1968.
- Selwood, A., Johnston, K., Katona, C., et al (2007) Systematic review of the effect of psychological interventions on family caregivers of people with dementia. *J Affect Disord*, 101, 75-89.
- Sitzer, D. I., Twamley, E. W. & Jeste, D. V. (2006) Cognitive training in Alzheimer's disease: a meta-analysis of the literature. *Acta Psychiatr Scand*, 114, 75-90.
- Skjerve, A., Bjorvatn, B. & Holsten, F. (2004) Light therapy for behavioural and psychological symptoms of dementia. *Int J Geriatr Psychiatry*, 19, 516-522.
- Smits, C. H., de Lange, J., Droles, R. M., et al (2007) Effects of combined intervention programmes for people with dementia living at home and their caregivers: a systematic review. *Int J Geriatr Psychiatry*, 22, 1181-1193.
- Snowden, M., Sato, K. & Roy-Byrne, P. (2003) Assessment and treatment of nursing home residents with depression or behavioral symptoms associated with dementia: a review of the literature. *J Am Geriatr Soc*, 51, 1305-1317.
- Staal, J. A., Sacks, A., Matheis, R., et al (2007) The effects of Snoezelen (multi-sensory behavior therapy) and psychiatric care on agitation, apathy, and activities of daily living in dementia patients on a short term geriatric psychiatric inpatient unit. *Int J Psychiatry Med*, 37, 357-370.
- Sung, H. C. & Chang, A. M. (2005) Use of preferred music to decrease agitated behaviours in older people with dementia: a review of the literature. *J Clin Nurs*, 14, 1133-1140.
- Sung, H. C., Chang, A. M. & Lee, W. L. (2010) A preferred music listening intervention to reduce anxiety in older adults with dementia in nursing homes. *J Clin Nurs*, 19, 1056-1064.
- Thorgrimsen, L., Spector, A., Wiles, A., et al (2003) Aroma therapy for dementia. *Cochrane Database Syst Rev*, CD003150.
- Vink, A. C., Birks, J. S., Bruinsma, M. S., et al (2004) Music therapy for people with dementia. *Cochrane Database Syst Rev*, CD003477.
- Voigt-Radloff, S., Graff, M., Leonhart, R., et al (2009) WHEDA study: effectiveness of occupational therapy at home for older people with dementia and their caregivers--the design of a pragmatic randomised controlled trial evaluating a Dutch programme in seven German centres. *BMC Geriatr*, 9, 44.
- Woods, B., Spector, A., Jones, C., et al (2005) Reminiscence therapy for dementia. *Cochrane Database Syst Rev*, CD001120.
- Woods, R. T., Bruce, E., Edwards, R. T., et al (2009) Reminiscence groups for people with dementia and their family carers: pragmatic eight-centre randomised trial of joint reminiscence and maintenance versus usual treatment: a protocol. *Trials*, 10, 64.
- Yamaguchi, H., Maki, Y. & Yamagami, T. (2010) Overview of non-pharmacological intervention for dementia and principles of brain-activating rehabilitation. *Psychogeriatrics*, 10, 206-213.