

Bewegungsanalyse für die Rehabilitation

Aktueller Stand und klinische Perspektiven

J. Steinhardt, A. Zolal, T. Heß, F. J. Marin, M. Kirsch

Literatur

- [1] Lam, W.W.T., Tang, Y.M. & Fong, K.N.K. A systematic review of the applications of markerless motion capture (MMC) technology for clinical measurement in rehabilitation. *J NeuroEngineering Rehabil* 20, 57 (2023). doi: 10.1186/s12984-023-01186-9
- [2] Steffensen EA, Magalhães F, Knarr BA, Kingston DC. Comparison of markerless and marker-based motion capture of gait kinematics in individuals with cerebral palsy and chronic stroke: A case study series. *Res Sq* [Preprint]. 2023 Feb 8;rs.3.rs-2557403. doi: 10.21203/rs.3.rs-2557403/v1
- [3] Zhou, Y., Rashid, F. 'A. N., Mat Daud, M., Hasan, M. K., & Chen, W. (2025). Machine Learning-Based Computer Vision for Depth Camera-Based Physiotherapy Movement Assessment: A Systematic Review. *Sensors*, 25(5), 1586. <https://doi.org/10.3390/s25051586>
- [4] Hou K, Li Y, Wang R, et al. (2025): An accessible AI-assisted rehabilitation system for real-time pose estimation and movement analysis. *Sensors*. 2025;25(19):6239. doi: 10.3390/s25196239
- [5] Armand S, Sawacha Z, Goudriaan M, Horsak B, van der Krogt M, Huenaerts C, Daly C, Kranzl A, Boehm H, Petrarca M, Guiotto A, Merlo A, Spolaor F, Campanini I, Cosma M, Hallemans A, Horemans H, Gasq D, Moissenet F, Assi A, Sangeux M. Current practices in clinical gait analysis in Europe: A comprehensive survey-based study from the European society for movement analysis in adults and children (ESMAC) standard initiative. *Gait Posture*. 2024 Jun;111:65-74. doi: 10.1016/j.gaitpost.2024.04.014.
- [6] Martínez-Cid S, Herrera V, Schez-Sobrinho S, et al. (2025): Artificial Intelligence in Physical Therapy for Neurological Rehabilitation: A Systematic Mapping Study. *IEEE Access*. 2025;13:1850655-1850785. doi: 10.1109/ACCESS.2025.3626013
- [7] Liu H, Xie Q. Applications of artificial intelligence in rehabilitation: technological innovation and transformation of clinical practice. *SLAS Technology*. 2025;35:100360. doi: 10.1016/j.slas.2025.10036
- [8] Alshami A, Nashwan A, AlDardour A, Qusini A. (2025): Artificial Intelligence in rehabilitation: a narrative review on advancing patient care. *Rehabilitación (Madr)*. 2025;59(2):100911. doi: 10.1016/j.rh.2025.100911
- [9] Minea, M., Ismail, S., Petcu, L. C., Nedelcu, A.-D., Petcu, A., Minea, A.-E., & Iliescu, M.-G. (2025). Using Computerised Gait Analysis to Assess Changes After Rehabilitation in Knee Osteoarthritis: A Systematic Review and Meta-Analysis of Gait Speed Improvement. *Medicina*, 61(9), 1540. <https://doi.org/10.3390/medicina61091540>
- [10] Sumner J, Lim HW, Chong LS, Bunde A, Mukhopadhyay A, Kayambu G. (2023): Artificial intelligence in physical rehabilitation: a systematic review. *Artificial Intelligence in Medicine*. 2023;146:102693. doi: 10.1016/j.artmed.2023.102693

- [11] Luo Z, Wang Y, Zhang T and Wang J (2025) Effectiveness of AI-assisted rehabilitation for musculoskeletal disorders: a network meta-analysis of pain, range of motion, and functional outcomes. *Front. Bioeng. Biotechnol.* 13:1660524. doi: 10.3389/fbioe.2025.1660524
- [12] Roggio F, Trovato B, Sortino M, Musumeci G. A comprehensive analysis of the machine learning pose estimation models used in human movement and posture analyses: A narrative review. *Heliyon*. 2024 Oct 30;10(21):e39977. doi: 10.1016/j.heliyon.2024.e39977.
- [13] Heitzer, F., Raab, D., Kowalczyk, W. et al. Orthopädie der Zukunft. *Orthopädie* 54, 991–1002 (2025). doi:10.1007/s00132-025-04708-6
- [14] Del Din S, Elshehabi M, Galna B, et al. Gait analysis with wearables predicts conversion to Parkinson disease. *Ann Neurol* 2019;86(3): 357–367. doi: 10.1002/ana.25548
- [15] Darweesh SKL, Verlinden VJA, Stricker BH, Hofman A, Koudstaal PJ, Ikram MA. Trajectories of prediagnostic functioning in Parkinson's disease. *Brain* 2017;140(2):429–441. doi: 10.1093/brain/aww291
- [16] Ishaq, M., Guastella, D. C., Sutera, G., & Muscato, G. (2026). A Systematic Review of Fall Detection and Prediction Technologies for Older Adults: An Analysis of Sensor Modalities and Computational Models. *Applied Sciences*, 16(4), 1929. doi:10.3390/app16041929
- [17] Putzolu M, Manzini V, Gambaro M, et al. (2023): Home-based exercise training using a smartphone app in patients with Parkinson's disease: a feasibility study. *Frontiers in Neurology*. 2023;14:1205386. doi: 10.3389/fneur.2023.1205386
- [18] Wagner L, Hauptmann B, Hoffmann AK, et al. (2022): Evaluation of an individualized, tablet-based physiotherapy training programme for patients with Parkinson's disease: the ParkProTrain study. *BMC Neurology*. 2022;22(1):176. doi: 10.1186/s12883-022-02647-9
- [19] Steinhardt J, Hanssen H, Heldmann M, Sprenger A, Domingo A, Reyes C, Rosales R, Klein C, Münte T, Westenberger A, Oropilla JQ, Diesta C, Brüggemann N. (2023): Prodromal phase of X-linked Dystonia-Parkinsonism is characterized by a subclinical motor phenotype. *Movement Disorders*. 2023. doi: 10.1002/mds.29033